



Getting Ready for Your Colonoscopy

2-Day GoLYTELY Prep

Physician's Name: _____

Procedure Date: _____

The time of your colonoscopy will be given to you three days before your appointment.

☐ **David H. Koch Center**

Endoscopy Center
1283 York Avenue, 8th Floor
New York, NY 10065

☐ **NewYork-Presbyterian
Hospital**

Central Endoscopy Suite
525 East 68th Street, 2nd Floor
(Greenberg Pavilion)
New York, NY 10065

☐ **NewYork-Presbyterian
Lower Manhattan Hospital**

Endoscopy Suite
170 William Street, 2nd Floor
New York, NY 10038

Why a Clean Colon Matters

Food and liquid can stay in your body for many hours. If your colon is not completely empty during your colonoscopy, your doctor may not be able to see **polyps or other findings**, and the exam may need to be rescheduled. Following these instructions carefully helps make sure your colon is empty when you arrive.

How to Contact Us

- **Questions about your colonoscopy or preparation:** Call **646-962-4000**, Monday–Friday (excluding holidays), 9:00 AM – 5:00 PM.
- **After 5:00 PM, weekends, or holidays:** An on-call physician will return your call. Non-urgent calls are returned the next business day.
- **To reschedule:** Call **646-962-4000**.
- **Medical emergency:** Call **911** or go to the nearest Emergency Room.

Cancellation Policy

Scheduling your colonoscopy requires careful planning between staff, doctors, and the hospital. To help avoid delays, please follow this guide and the schedule.

If you must reschedule or cancel:

Contact us **at least 3 days before** your colonoscopy.

Phone: 646-962-4000 **Email:** endoscopyquestions@med.cornell.edu

Contents

2-Day GoLYTELY Prep

Getting Ready for Your Colonoscopy	4
Cardiac Clearance	6
Medication Information	6
Anticoagulants & Antiplatelets	6
Diabetes & Weight-Loss Medications	7
Blood Pressure Medications	7
Iron Supplements	7
What to Buy to Clean Your Colon	8
Eating Before Your Colonoscopy	9
What is a Low-Fiber Diet?	9
What is a Clear-Liquid Diet?	10
2-Day GoLYTELY Prep Instructions	11
What to Expect	12
Day of Your Colonoscopy	12
Before Your Colonoscopy	12
In the Recovery Room	12
At Home, After Your Colonoscopy	12

Getting Ready for Your Colonoscopy

As soon as your colonoscopy is scheduled:

- Obtain cardiac clearance, if applicable (see page 6).
- Review medication information for medicines you may need to stop (see pages 6 and 7).
- Buy the products you need to clean your colon (see page 8).

7 days before your colonoscopy (see pages 7 and 9):

- Take 1 capful of MiraLAX daily.
- Stop iron supplements.
- Stop weekly GLP-1 receptor agonists.
- Begin a low-fiber diet.

3 days before your colonoscopy (see page 7):

- Stop SGLT2 inhibitors.

2 days before your colonoscopy (see pages 9, 10, and 11):

- Clear liquids all day — no solid food.
- 9:00 AM — Mix the first bottle of GoLYTELY by adding warm water up to the fill line. Refrigerate.
- 5:30 PM — Drink an 8-oz glass of GoLYTELY every 10–15 minutes until you finish HALF of the bottle. Refrigerate the remainder.

1 day before your colonoscopy (see pages 9, 10, and 11):

- Clear liquids all day — no solid food.
- 9:00 AM — Mix the second bottle of GoLYTELY by adding warm water up to the fill line. Refrigerate.
- 12:00 PM — Take 2 Dulcolax tablets.
- 5:00 PM — Drink an 8-oz glass of GoLYTELY every 15 minutes until you finish the entire second bottle.

Day of your colonoscopy (see page 12):

5 hours before your arrival time:

- Take one 250 mg or two 125 mg Gas-X tablet(s).
- Drink the remaining HALF of the first bottle of GoLYTELY, an 8-oz glass every 15 minutes. Finish within 2 hours.
- Drinking the prep 5 hours before your arrival time helps your doctor see polyps.

3 hours before your arrival time:

- Do not drink or eat anything — including water, any remaining preparation, mints, candies, or gum.

Arrival Time Reference

Use the row that matches your scheduled arrival time. Your final procedure time will be confirmed three days in advance.

Arrival Time	Start drinking your remaining prep	Finish drinking the entire prep by
6 AM	1 AM	3 AM
7 AM	2 AM	4 AM
8 AM	3 AM	5 AM
9 AM	4 AM	6 AM
10 AM	5 AM	7 AM
11 AM	6 AM	8 AM
12 PM	7 AM	9 AM
1 PM	8 AM	10 AM
2 PM	9 AM	11 AM

Cardiac Clearance

If you have a cardiac defibrillator (AICD), permanent pacemaker (PPM), or any current cardiac problems, we need a letter from your cardiologist **at least 1 week before your colonoscopy**. Cardiac problems include any history of heart attack, chest pain, difficulty breathing, or fainting episodes.

If you have...	We need...
Permanent Pacemaker (PPM)	An interrogation report within 12 months
Cardiac Defibrillator (AICD)	An interrogation report within 6 months

Medication Information

Anticoagulants (Blood Thinners)

If you take a blood thinner, **ask the doctor who prescribed it** whether you should stop it before your colonoscopy. If your prescribing doctor does not want you to stop the medication, please call our office.

Commonly prescribed medications:

Apixaban (Eliquis®)	Edoxaban (Savaysa®)	Ticagrelor (Brilinta®)
Cilostazol (Pletal®)	Enoxaparin (Lovenox®)	Tinzaparin (Innohep®)
Clopidogrel (Plavix®)	Heparin	Warfarin (Coumadin®)
Dabigatran (Pradaxa®)	Prasugrel (Effient®)	Fondaparinux (Arixtra®)
Dalteparin (Fragmin®)	Rivaroxaban (Xarelto®)	Ticlopidine (Ticlid®)

DO NOT STOP YOUR ASPIRIN unless your doctor specifically tells you to do so.

Diabetes & Weight-Loss Medications

If you take...	Do this
Insulin or oral hypoglycemic pills	Check with the doctor who prescribed these for you, as soon as possible, for instructions on how to manage them around your colonoscopy.
SGLT2 inhibitors Canagliflozin (Invokana, Invokamet); Dapagliflozin (Farxiga, Xigduo XR, Qtern); Empagliflozin (Jardiance, Synjardy, Glyxambi, Trijardy XR); Ertugliflozin (Steglatro, Segluromet, Steglujan); Sotagliflozin (Inpefa); Bexagliflozin (Brenzavvy)	Stop 3 days before your colonoscopy.
Daily GLP-1 receptor agonists Liraglutide (Victoza, Saxenda, Xultophy); Exenatide (Byetta); Lixisenatide (Adlyxin, Soliqua); Semaglutide oral tablet (Rybelsus, Wegovy); Orforglipron (Foundayo)	Do not take your dose on the day of your colonoscopy. You will also need to take MiraLAX every day for 1 week before your colonoscopy (see page 12).
Weekly GLP-1 receptor agonists Dulaglutide (Trulicity); Exenatide ER (Bydureon BCise); Semaglutide injection (Ozempic, Wegovy); Tirzepatide (Mounjaro, Zepbound)	Stop 1 week before your colonoscopy. You will also need to take MiraLAX every day for 1 week before your colonoscopy (see page 12).

Blood Pressure Medications

Take your blood pressure medications as usual on the day of your colonoscopy with a small sip of water — **except diuretics** (water pills). **Do not take diuretics** on the morning of your colonoscopy.

Iron Supplements

Stop oral iron supplements **7 days before your exam**. You may resume them after your colonoscopy.

What to Buy to Clean Your Colon

Your bowel prep requires a prescription from your doctor. All other items below are available over the counter — no prescription needed.

Product	Quantity	Where to Buy	What it looks like
GoLYTELY <i>or Gavilyte, NuLYTELY, or generic polyethylene glycol electrolyte solution</i>	2 bottles	Pharmacy (prescription required)	
Dulcolax (bisacodyl) <i>or generic bisacodyl — NOT the stool softener</i>	2 tablets	Pharmacy	
Gas-X (simethicone) <i>or generic simethicone</i>	Two 125 mg or one 250 mg tablet	Pharmacy	
MiraLAX <i>or generic polyethylene glycol 3350</i>	1 bottle (238 g)	Pharmacy	

Eating Before Your Colonoscopy

Seven Days Before Your Exam

Begin a low-fiber diet starting one week before your exam.

Two Days Before Your Exam

Clear liquids all day. Do not eat any solid food.

The Day Before Your Exam

Clear liquids all day. Do not eat any solid food.

What is a Low-Fiber Diet?



Acceptable Foods:

Chicken, fish, beef, pork, tofu, eggs, white rice, white bread, plain bagel, white pastas or noodles, cooked potatoes without the skin, bananas, avocado, vanilla ice cream, plain yogurt, cheese, and butter.

Foods to Avoid:

Nuts, seeds, corn, beans, quinoa, raw fruits and vegetables, skin of cooked fruits and vegetables, whole-grain or whole-wheat products. These foods are hard to digest, do not easily clear out of your colon, and make it difficult for your doctor to see your colon clearly during the procedure.

What is a Clear-Liquid Diet?

Chicken Broth or Bouillon



Jell-O or Popsicles



Ginger Ale or Clear Juice



Acceptable clear liquids:

Water; black coffee or tea (no milk or creamer); ginger ale or other clear sodas; sports drinks; clear broth or bouillon (no noodles or solids); clear juices (apple, white grape); plain Jell-O; frozen juice popsicles.

Avoid:

Any red or purple liquids; alcohol; juice with pulp; milk, cream, or non-dairy/soy creamer; soup with solids; smoothies; or any liquid you cannot see through.

2-Day GoLYTELY Prep Instructions

One Week Before Your Exam

- Mix 1 capful (17 g) of MiraLAX in an 8-oz beverage and drink once daily, every day, for the week before your colonoscopy.
- Begin a low-fiber diet.

Two Days Before Your Exam

- 9:00 AM — Mix the first bottle of GoLYTELY by adding warm water up to the fill line. Refrigerate.
- 5:30 PM — Drink an 8-oz glass of GoLYTELY every 10–15 minutes until you finish HALF of the first bottle. Refrigerate the remainder to drink on the day of your colonoscopy.

One Day Before Your Exam

- 9:00 AM — Mix the second bottle of GoLYTELY by adding warm water up to the fill line. Refrigerate.
- 12:00 PM — Take 2 Dulcolax tablets with an 8-oz glass of water or other clear liquid.

One Day Before Your Exam, 5:00 PM

- Drink an 8-oz glass of GoLYTELY every 15 minutes until you finish the entire second bottle.

What to expect:

You will have loose, frequent stools — they may not start until after the next step. Use soft toilet paper, and apply petroleum jelly to the perianal area at the start of the diarrhea to reduce irritation.

Continue drinking even if you have diarrhea. If you feel nauseated, take a 60-minute break, then resume.

5 Hours Before Your Arrival Time

- Take one 250 mg or two 125 mg Gas-X tablet(s).
- Drink the remaining HALF of the first bottle of GoLYTELY, an 8-oz glass every 15 minutes. Finish within 2 hours.

It is important to finish the entire prep so the colon is fully cleansed. If you feel nauseated, pause until it resolves and then continue.

3 Hours Before Your Arrival Time

- **STOP** any remaining preparation.
- Do not drink or eat anything — including water, mints, candies, or gum.
- **If you eat or drink anything during this window, your colonoscopy must be cancelled.**

You may take your usual medications — unless instructed otherwise — with a small sip of water.

What to Expect

Day of Your Colonoscopy

A colonoscopy is a medical procedure in which your doctor examines the colon and parts of the small intestine using a flexible scope with a camera. During the procedure, your doctor will look at the colon walls to check for any findings that may need treatment, including polyps.

IMPORTANT — Escort Required

Per hospital policy, an adult age 18 or older must pick you up and escort you home after your procedure.

- The procedure cannot begin until this plan is confirmed. If unconfirmed, your appointment will be cancelled.
- Plan for pickup approximately 1.5 to 2 hours after your scheduled procedure time.
- If you don't have an escort, please let us know — we can provide information on service agencies that will accompany you (at cost).

Before Your Colonoscopy

When you arrive, you will register and sign forms. You may also check in online through Connect, the patient portal for NewYork-Presbyterian, Columbia, and Weill Cornell Medicine. You will be asked to state and spell your name and date of birth to verify your identification.

After changing into a gown, a nurse will place an intravenous (IV) catheter in your arm. An anesthesiologist will review your medical history. Your doctor will give you more information about the procedure, including any risks, and answer your questions.

In the procedure room, you will be connected to monitors for your heart, breathing, and blood pressure, and given oxygen through your nose. Anesthesia is administered through your IV. Your doctor will use an endoscope to examine your colon. If polyps are found, the doctor will remove them and send them to the laboratory. A colonoscopy is generally not painful.

In the Recovery Room

You will wake up in the recovery room. Some bloating or stomach cramping is normal and resolves with passing gas.

Once you are fully awake, your nurse will remove the IV. Your doctor will then review your colonoscopy results with you. Your nurse will go over discharge instructions before you leave. Your escort will take you home.

At Home, After Your Colonoscopy

If you had a biopsy, you may notice a few drops of blood from the rectum. This is normal after a biopsy. Avoid alcohol for 24 hours. Avoid carbonated drinks, raw fruit, and salads at your first meal. You may resume your normal diet and activities the next day.

Call your doctor's office if you have:

- A temperature of 101°F (38.3°C) or higher
- Severe abdominal pain, or increasing abdominal size or firmness
- Heavy bleeding from the rectum lasting more than 24 hours
- Weakness, faintness, or persistent nausea or vomiting